

Walter C. Pettus
Yale University
(formerly UW–Madison)

SSGF Annual Program Review
29 June – 2 July 2015

Background Characterization and Detector Sensitivity in DM-Ice

Understanding the Universe

Visible Sky



Axel Mellinger
http://apod.nasa.gov/apod/ap010202.html

Understanding the Universe

Gamma Ray Sky



Fermi LAT Collaboration
http://fermi.gsfc.nasa.gov/ssc/Fermi_5_year.jpg

Understanding the Universe

Microwave Sky



Planck Collaboration
arXiv:1502.01582

Understanding the Universe

Microwave Sky



Planck Collaboration
arXiv:1502.01582

A Dark Universe






Millennium Run
10.077.696.000 particles


cz / H0DL
redshift z
0.01
0.1
1
0.5
1
Ωm=1
Ωm=0.3
Ωm=0.3 k=0
Ωm=0
Ωm=0 k=0




Dark Matter
26.8%
Ordinary Matter
4.9%
Dark Energy
68.3%


baryon density Ωbh2
10-2
Y
D/H
3He/H
7Li/H
CMB
BBN


Planck
WMAP9
ACT
SPT
Dℓ [μK2]
ℓ




V (km s-1)
R(kpc)

Detecting Dark Matter

Indirect Searches

- Identify annihilation/ decay products of dark matter from extraterrestrial sources





Collider Searches

- Identify production signatures of dark matter in collision



Direct Searches

- Identify energy deposition from dark matter interaction in a terrestrial detector



Experiment Design – DAMA

Monolithic detectors
(up to ton-scale)

Component radiopurity
(control of radioactive backgrounds)

Environmental stability and control
(temperature, radon, …)

Performed search for dark matter annual modulation with NaI(Tl) crystal array



Bernabei *et al.*, NIM A, (2008)

Experiment Design – DAMA

Monolithic detectors
(up to ton-scale)

Component radiopurity
(control of radioactive backgrounds)

Environmental stability and control
(temperature, radon, …)

Low-energy (~1 keV) thresholds necessary for dark matter sensitivity
(characterization and removal of noise)

Performed search for dark matter annual modulation with NaI(Tl) crystal array



Bernabei *et al.*, NIM A, (2008)

Demonstrated excellent background reduction:



Bernabei *et al*, Eur Phys J C (2008)

DAMA/LIBRA – Detection?

A Solitary (and Persistent) Signal

- Observed modulating signal with ***9.3-sigma*** significance
- Matches dark matter expectation



Bernabei *et al*, arXiv:1412.6524

Performed search for dark matter annual modulation with NaI(Tl) crystal array



Bernabei *et al.*, NIM A, (2008)



Demonstrated excellent background reduction:



Bernabei *et al*, Eur Phys J C (2008)

Tension of Experimental Results

A Solitary (and Persistent) Signal

- Observed modulating signal with ***9.3-sigma*** significance
- Matches dark matter expectation



Bernabei *et al*, arXiv:1412.6524

Conflict with Null Results

- DAMA remains the only experiment to claim dark matter detection
- Strongly excluded by numerous other experiments



PandaX Collaboration, arXiv:1505.00771

Resolution hidden in
Astrophysics?
Particle Physics?
Instrumental Effects?
Uncharacterized Background?

DM-Ice Experiments

DM-Ice17

(*January 2011 – present*)



First dark matter experiment in South Pole ice

- Demonstrated viability and advantage of environment

DM-Ice37

(*April 2014 – present*)



R&D testbed for NaI detectors

- Crystal background
- Light yield
- PMT/lightguide configurations

DM-Ice250

(*future*)



Science result

- Definitive test of DAMA dark matter claim

See also: Cherwinka *et al.*, Astropart. Phys **35** (2012)

A World of Dark Matter Searches



A World of Dark Matter Searches



DM-Ice17 Experiment

(2x) 8.5-kg NaI(Tl) modules

- Installed Dec 2010
- Data run from June 2011 to January 2015

Goals:

- Demonstrate the feasibility of deploying and operating NaI(Tl) detectors in the Antarctic Ice for a dark matter search
- Study environmental stability
- *In situ* measurement of the radiopurity of the Antarctic ice / hole ice at 2450 m depth
- Study the capability of IceCube to veto muons

First results published:

Cherwinka *et al.*, Phys. Rev. D **90** (2014)





South Pole Science

Geographic South Pole



Amundsen-Scott South Pole Station

Skiway



DM-Ice17

IceCube
IceTop

Keck Array

BICEP3, SPT

South Pole Logistics



Walter C. Pettus

DM-Ice17 Uptime

> 99% uptime



Sources of downtime understood (DAQ, characterization data, satellite issues)

- Uptime has improved over physics data run



Temperature Stability

- Ice environment provides stable temperature
  - 0.3°C cooling over 3.5 years measured at mainboard
  - < 0.025°C daily RMS

Changing PMT HV setting affects MB temperature

Power outages affect MB temp, not crystal temp



DM-Ice17 Background

Betas & Gammas:

- Internal contamination lines allow calibration
- Well matched by *a priori* background model
  - Only crystal background unconstrained

Alphas

- Crystal contaminants only
- Constrains ^{232}Th-, ^{235}U-, ^{238}U- and chain levels





Detector backgrounds dominate at all energies;
Negligible contributions from environmental background

Cherwinka *et al.*, Phys. Rev. D **90** (2014)

Cosmic Ray Air Showers



Louis *et al.*, Los Alamos Science (1997)

High energy particles, 'cosmic rays' penetrating through Earth's atmosphere produce cascades of secondary particles following interaction

- cosmic ray intensity varies with position on Earth
- shower intensity varies with altitude
- secondary particles are a source of both prompt and delayed background

Cosmic Ray Air Showers



* shower greatly simplified

10 TeV Proton Shower
(millions of secondaries)

CORSIKA Simulation, https://www.ikp.kit.edu/corsika/



Louis *et al.*, Los Alamos Science (1997)

Prompt Cosmic Background



In preparation, under collaboration review

Cosmic ray muons passing directly through DM-Ice detectors

- Highest energy signals measured
  - Unambiguous identification of muons with IceCube detector demonstrated
- Induces seconds-long afterpulsing



Delayed Cosmic Background

^{127}I (n,3n) ^{125}I



Cosmic ray hadrons interact with nuclei in detector components

- Transmutation of nucleus to radioisotope via spallation or capture processes
- Decay over days or years governed by radioactive half-life

Delayed Cosmic Background

^{127}I (n,3n) ^{125}I



Cosmic ray hadrons interact with nuclei in detector components

- Transmutation of nucleus to radioisotope via spallation or capture processes
- Decay over days or years governed by radioactive half-life

- Neutrons dominate cosmogenic activation rates
- Activation cross section peaked for 10 – 100 MeV neutrons
- Activation rates scale with cosmic ray rate
  - Dependence on altitude, position, …

Position Scaling



Lifton *et al.*, Earth and Plan. Sci Lett. (2014)

Cosmogenic Activation Hazards

Location	Relative Neutron Rate (to sea level)
Madison, WI	1.38
South Pole	11.1
Commercial Flight	100 – 600



Cosmic Ray Exposure Timeline

Full Detector Component History



F_{tot} – Relative cosmic ray neutron flux (scaling from sea level)

- Long periods of low-level exposure during storage and construction
- Punctuated exposure from flight shipment

"Cumulative Activation" – Time-integrated neutron flux scaling

- Different detector components have different exposure histories
- Two DM-Ice17 detectors have different deployment times

Cosmic Ray Exposure Timeline

Full Detector Component History



Flight Segments



But decay not included: $\lambda N = R \cdot \left(1 - e^{-\lambda t}\right)$

Cosmogenic Prediction

- Most cosmogenic features arise from NaI activation
  - Rates simulated for start of dataset (6 months of decay)
  - Five peaks (four isotopes) with > 1 dru rate
    - All of which have been identified in data





Cosmogenic Decay Peaks



Confirm identity of cosmogenic peaks:

- Match simulated spectral features
  - Expect 65.3 keV full-energy and 37.6 keV L-shell capture peaks
- Measuring decay time
  - Expect 59.4 day half-life



Low-Energy Spectrum

Simulation can reproduce changing spectral features

- Requires 6 cosmogenic isotopes
- Improved total background model via broken chains (^{232}Th in pressure vessel)
- Independent measurement of ^{60}Co ($t_{1/2}$ = 5.3 yr)
- Improvement in energy resolution modeling
  - Mismatch of 65 keV peak width
- Region of interest rates constrained indirectly
  - All 3 keV cosmogenic components have higher-energy peaks



DM-Ice17 Cosmogenics

- Good agreement with other NaI activation measurements
  - Validation of scaling model
- Identification of one new activated radioisotope (^{113}Sn)
  - Implications for low-energy region of interest

	Isotope	$t_{1/2}$ (days)	DM-Ice17 (#/kg/day)	Literature (#/kg/day)
requires multi-crystal →	^{22}Na	951	-	45.1 ± 1.9
	^{113}Sn	115.1	16	-
short half-life →	^{121}Te	19.2	-	9.9 ± 3.7
	^{121m}Te	164.2	25	23.5 ± 0.8
discrepant →	^{123m}Te	119.2	21	31.6 ± 1.1
	^{125m}Te	57.4	27	28.2 ± 1.3
	^{127m}Te	106.1	< 9	10.2 ± 0.4
	^{125}I	59.4	230	220 ± 10
short half-life →	^{126}I	12.9	-	283 ± 36

DM-Ice250 Experimental Program

Modular detector supporting deployment in both hemispheres



- 125 kg NaI(Tl) / module
- 7-crystal array
  - Coincidence between PMTs
  - Anti-coincidence between crystals

DM-Ice250N



DM-Ice250S



Background Simulation

- Crystal array geometry allows for coincidence studies
  - Reduction in total background rate via anti-coincidence
  - Stronger constraint on background model via MC comparison
- Crystal backgrounds dominate at all energies

Outer Crystal Spectrum
(*after veto*)

- ^{40}K 3 keV peak complicates ROI
  - Veto reduction of 37-67%
- ^{210}Pb flat background is remaining low-energy background
  - Minimal improvement with veto
- High light yield imperative for resolution and noise reduction
  - 15 pe/keV demonstrated in alternate geometry



DM-Ice250S Activation

relative to DM-Ice17:



Maintain polar program flights

Minimize storage time



DM-Ice250S Cosmogenics

Event rate one month after deployment

- Multiple strong cosmogenic calibration lines
- Significant contributions to 2 – 6 keV region of interest

Cosmogenic contribution to ROI:

- ^{126}I ($t_{1/2}$ = 13 days)
  - lead contribution at deployment
- ^{113}Sn ($t_{1/2}$ = 115 days)
  - dominates rate over physics run





Cosmogenic Mitigation

"Exposure budget" for ^{113}Sn in DM-Ice250S:

- After "easy" 40% reduction
- Major contributions remain from NZL-McM flight and South Pole



Further reductions:

- 50% reduction in low-altitude NZL-McM cargo flight (15% of total)



- 90% reduction in South Pole activation from under-ice storage



Sensitivity Projection

Null Hypothesis:
(90% CL contours, 500 kg*yr exposure)



Signal Hypothesis:
(250 kg detector)



DAMA dark matter claim can be definitively tested with 500 kg*yr exposure

- 5-sigma test possible with background of 2 dru
  - *cf.* DAMA background of ~1 dru



- DM-ICE17 has completed a successful 3.5-yr data run
  - 1 concept paper and 1 results paper published
  - +4 more papers by year end
- NaI(Tl) continues to have a promising future as a detector for dark matter searches
  - Verification or refutation of DAMA within few years
  - Still learning about activation products, phosphorescence,…
  - DM-ICE250 uniquely poised as planned Southern Hemisphere experiment
- Four detector built and operated
  - 3 NaI(Tl) and 1 plastic scintillator
    - (last R&D detector has been deconstructed and rebuilt 2 more times)
  - Uncountably many tons of lead moved

Some parting thanks:

To the NNSA for their sponsorship of this fellowship program

To Krell for four great years of support and coordination

To you for stimulating discussions at these conferences



LUX ET VERITAS

WIPAC
WISCONSIN ICECUBE PARTICLE ASTROPHYSICS CENTER

SSGF
STEWARDSHIP SCIENCE GRADUATE FELLOWSHIP

THE UNIVERSITY of WISCONSIN MADISON